

# Similarity properties and scaling laws of two-temperature radiating fluids

E. Falize<sup>1</sup>, A. Dizière<sup>2</sup>, B. Loupiau<sup>1</sup>, M. Koenig<sup>2</sup>

<sup>1</sup>*CEA-DAM-DIF, F-91297 Arpajon, France, [Emeric.Falize@obspm.fr](mailto:Emeric.Falize@obspm.fr)*

<sup>2</sup>*LULI, Ecole Polytechnique, CNRS, CEA, UPMC, Route de Saclay, 91128 Palaiseau, France*

Several Astrophysics accretion environments, such as accreted column in magnetic cataclysmic variables [1], are such as the radiative cooling time-scale of the plasma is shorter than the electron-ion collision time-scale. Consequently the electrons will lose their energy rapidly while their energy gains by collisions are unable to keep up radiative loss [2,3]. In this case the two-temperature effects are significant which considerably complicate the structure and the emission of plasma [1,4]. In this work we discuss the specificity of this regime, the scalability and the self-similarity behaviour of flows in such radiating regime. This analyze complete the precedent studies that they have been realized [5,6].

## References

- [1] Imamura J. N. et al., *Astrophys. J.* 458: 327 (1996)
- [2] Zeldovich Y. B. & Raizer Y. P., *Physics of Shock Waves and High-Temperature hydrodynamic phenomena* (Academic, New York, 1966)
- [3] Kylafis N. D. & Lamb D. Q., *Astrophys. J.* 48: 239 (1982)
- [4] Saxton C. J. et al., *MNRAS* 360: 1091 (2005)
- [5] Falize E. et al., *Astrophys. Spac. Sci.* 322: 107 (2009)
- [6] Falize E. et al. *Astrophys. J.* (submitted), ArXiv 0910.2338F